

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED		1b. RESTRICTIVE MARKINGS NONE	
2a. SECURITY CLASSIFICATION AUTHORITY DECLASSIFICATION/DOWNGRADING SCHEDULE		3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release. Distribution unlimited.	
4. PERFORMING ORGANIZATION REPORT NUMBER(S) Technical Report No. 15		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
NAME OF PERFORMING ORGANIZATION Massachusetts Institute of Technology	6a. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION ONR	
ADDRESS (City, State, and ZIP Code) 77 Massachusetts Avenue, Room 1-306 Cambridge, MA 02139		7b. ADDRESS (City, State, and ZIP Code) 800 North Quincy Street Arlington, VA 22217	
NAME OF FUNDING/SPONSORING ORGANIZATION DARPA	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER N00014-86-0768	
ADDRESS (City, State, and ZIP Code) 1400 Wilson Boulevard Arlington, VA 22209		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO. R & T Code	PROJECT NO. A 400005
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) MOLECULAR STRUCTURE OF AMORPHOUS BISPHENOL-A POLYCARBONATE			
12. PERSONAL AUTHOR(S) Michelle Hutnik, Ali S. Argon, and Ulrich W. Suter			
13a. TYPE OF REPORT Abstract	13b. TIME COVERED FROM 1988 TO 1989	14. DATE OF REPORT (Year, Month, Day) 1989, May 29	15. PAGE COUNT 2
16. SUPPLEMENTARY NOTATION Abstract of oral presentation to be made at the American Chemical Society Symposium on Computer Simulation of Polymers on September 11-14, 1989 at Miami Beach.			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP	SUB-GROUP	
19. ABSTRACT (Continue on reverse if necessary and identify by block number) A detailed atomistic molecular mechanics model has been developed for the polycarbonate of bisphenol-A (PC) and is employed here to generate dense (glassy) microstructures. The resulting microstructures are being used to investigate the inelastic behavior of PC. The phenylenering flip is a feature of PC which has been widely studied using various NMR techniques. It is the simplest of inelastic processes and is amenable for simulation and study. The energy barrier to the ring flip has been obtained using the minimized structures, and the calculated mean energy barrier is 10.4 (±6.0) Kcal/mole. This value agrees very well with the NMR results that have been reported. <i>Model Ring Rotation Computer and Simulation</i>			
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION Unclassified	
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. Kenneth Wynne		22b. TELEPHONE (Include Area Code) (202) 696-4410	22c. OFFICE SYMBOL

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted.

All other editions are obsolete

SECURITY CLASSIFICATION OF THIS PAGE

U.S. Government Printing Office: 1985-507-047

(To be filled in by Division)

A. DIVISION OF Polymer Chemistry
B. TITLE OF PAPER Molecular Structure of Amorphous Bisphenol-4 Polycarbonate

Paper number as listed on program _____

Time Required for Presentation _____

☐ Poster Presentation Preferred

C. AUTHORS
Underline name of speaker

M. Hutnik
A.S. Argon
U.W. Suter

D. Business Mailing Address Including Zip Code and telephone Number
List Address only once if all authors at same address.
 Massachusetts Institute of Technology
 77 Massachusetts Ave.
 Room 1-306
 Cambridge, MA 02139

E. ACS Division **F. American Chemist**
 Member? Member?
☒ Yes ☒ Yes
☐ No ☐ No

Engineer?
If not, give classification such as biologist, physicist, etc. Ph.D?

☐ Chemist
☐ Chemical Engineer

Other Materials Scientist

NOTE: All presenting authors must register for the meeting —either full meeting registration or one-day registration for the day of presentation.

G. Work done at M. I. T.

H. Plan ACS nonACS Yes publication. Where? Undecided No Uncertain

I. Specify Equipment Required for Presentation Other than 2" x 2" slide or overhead (transparency) projector none

J. ABSTRACT. Please be BRIEF—150 words maximum if possible. Title of paper should be ALL CAPS; author(s) listed by first name, middle initial, last name; indicate full address w/zip code. SINGLE SPACE, BLACK CARBON RIBBON.

DO NOT
 USE →

MOLECULAR STRUCTURE OF AMORPHOUS BISPHENOL-A
 POLYCARBONATE. M. Hutnik*, A. S. Argon*, and U. W. Suter**,
 (*)Massachusetts Institute of Technology Cambridge, MA 02139, (**)Institut fuer Polymere, ETH-Zuerich, CH-8092 Zuerich, Switzerland.

A detailed atomistic molecular mechanics model has been developed for the polycarbonate of bisphenol-A (PC) and is employed here to generate dense (glassy) microstructures. The resulting microstructures are being used to investigate the inelastic behavior of PC. The phenylene ring flip is a feature of PC which has been widely studied using various NMR techniques. It is the simplest of inelastic processes and is amenable for simulation and study. The energy barrier to the ring flip has been obtained using the minimized structures, and the calculated mean energy barrier is 10.4 (\pm 6.0) Kcal/mole. This value agrees very well with the NMR results that have been reported.



By _____	
Distribution/ _____	
Availability Codes _____	
Dist	Avail and/or Special
A-1	

DO NOT
 USE →

K. MAIL ABSTRACT TO PERSON NAMED IN ACS DIVISIONAL DEADLINES PUBLISHED (JUNE & DEC.) IN C&EN

MOLECULAR STRUCTURE OF AMORPHOUS BISPHENOL-A POLYCARBONATE

M. Hutnik*, A. S. Argon*, and U. W. Suter**

(*)Massachusetts Institute of Technology
Cambridge, MA 02139

and
(**)Institut fuer Polymere, ETH-Zuerich
CH-8092 Zuerich, Switzerland

INTRODUCTION

A detailed atomistic molecular mechanics model has been developed for the polycarbonate of bisphenol-A (PC). The technique used has been successfully applied earlier to simple vinyl polymers and is employed here to generate dense (glassy) packing of PC [1,2]. The resulting microstructures are being used to investigate the inelastic behavior of PC. The first such application is the phenylene ring flip which occurs in glassy PC. The ring flip is a feature of PC which has been widely studied using various NMR techniques [3,4] and as the simplest of inelastic processes is amenable for simulation and study using the generated microstructures.

MODEL

To be able to model PC, it is necessary to first understand the intramolecular interactions of the PC repeat unit. We developed a molecular mechanics force field of the PC repeat unit that is appropriate for the microstructure generation technique, based on fixed bond angles and bond lengths. A Lennard-Jones potential energy function was used to represent the Van der Waals interatomic interactions while the Coulombic interactions are represented as a truncated Block-Walker potential [2,5]. Intrinsic torsional potentials specific to PC were constructed and are included in the force field. The details of the force field will be described in full at a future time.

Amorphous PC microstructures were generated using the above force field and the modelling technique developed by Theodorou and Suter [1]. The model is static, with temperature only entering in the choice of the density, and includes spatially periodic continuation. The periodic cube edge length is 18.44 Angstroms, the degree of polymerization is 17, and the density is 1.20 g/cm³, which is the experimental value at 298 K.

Initial chain conformations of the microstructures were generated using a Monte Carlo technique with the rotational isomeric state scheme previously developed for PC [6]. A staged energy minimization procedure is then applied to the microstructure to obtain well relaxed, amorphous systems (Figure 1, hydrogens are omitted for clarity). No vestiges of crystallinity are observed. The energy minimization for PC is not straight forward, requiring approximately 30 stages in which the atomic radii and the partial point charges are gradually 'blown up'. The difficulty of this minimization is due to the size of the rigid moiety in the PC repeat unit. 12 minimized microstructures have been generated so far.

SIMULATION OF RING FLIP

The phenylene ring flip that occurs in PC has been a topic of intense experimental investigation using various NMR techniques [3,4]. The energy barrier of the ring flip is due to a combination

of both intra- and intermolecular interactions. Since glass densities are very high, the intermolecular contributions are high and could potentially dominate the energy barrier. Using the minimized PC microstructures it is possible to obtain this barrier and compare the results with the reported experimental values.

To calculate the energy barrier to ring flip, a phenylene ring is incrementally rotated and then 'fixed' while the rest of the structure is allowed to relax. This process is repeated until a 'flip' is performed in the laboratory reference frame. Every ring of one microstructure is treated so that a sufficient statistical sample is obtained and possible spatial effects can be studied.

RESULTS AND DISCUSSION

A typical rotation path is displayed in Figure 2. It shows the system potential energy vs. torsional angle (true rotation in parentheses) for the third ring along the chain. The true rotation is defined as the angle between the plane of the ring at the onset of the experiment and the plane of the ring at present. Figures 3 and 4 show the microstructure (hydrogens are omitted and the atomic radii are shrunk so that the rotation is easily visible) at two different points along the rotation path; Figure 3 contains the initial configuration of the chain (torsional angle=57°, true rotation=0°), and Figure 4 gives the configuration at the peak energy (torsional angle=117°, true rotation=55°).

In general for the ring flip, preceding the initial energy peak the conformational changes that occur as the structure relaxes are reversible, whereas the conformational changes are irreversible after the energy peaks. This corresponds to the unrestrained relaxation from a saddle point configuration in transition state theory.

The results of the ring flip calculations are as follows (value in parenthesis is standard deviation of the mean):

34 barriers calculated
mean energy barrier = 10.4 (±6.0) Kcal/mole

The frequency distribution of the calculated results of the ring flip are shown in Figure 5.

These results compare very favorably to the NMR results that have been calculated:

- Proton spin-lattice relaxation:
12 Kcal/mole [3]

- Dipolar Rotational Spin-Echo ¹³C NMR:
11 Kcal/mole [4]

ACKNOWLEDGEMENTS

We gratefully acknowledge financial support by DARPA and the Office of Naval Research under contract No. N00014-86-K-0768. We would also like to thank Drs. F. Gentile and P. J. Ludovice from ETH for many invigorating discussions.

REFERENCES

- (1) Theodorou, D. N.; Suter, U. W. Macromolecules 1985, 18, 1467.
- (2) Ludovice, P. J.; Suter, U.W. Polym. Preprints 1987, 28, 295.

- (3) Jones, A. A.; O'Gara, F. F.; Inglefield, P. T.; Bendler, J. T.; Yee, A. F.; Ngai, K.L. *Macromolecules* 1983, 16, 658.
- (4) Schaefer, J.; Stejskal, E.O.; McKay, R.A.; Dixon, W.T. *Macromolecules* 1984, 17, 1479.
- (5) Block, H.; Walker, S. M. *Chem. Phys. Let.* 1973, 19, 363.
- (6) Hutnik, M.; Suter, U. W. *Polym. Preprints* 1987, 28, 293.




Figure 1. Periodic cube of PC, hydrogens omitted for clarity.




Figure 2. Ring flip rotation path for the third ring along the chain.




Figure 3. Initial configuration of chain (hydrogens omitted and atomic radii reduced for clarity), torsional angle = 57°, true rotation = 0°.




Figure 4. Configuration at the energy peak, torsional angle = 117°, true rotation = 55°.




Figure 5. Frequency distribution of the energy barriers.